



FOWL CHOLERA VACCINE, INACTIVATED

DIRECTIONS FOR USE

DESCRIPTION

Fowl Cholera Vaccine, Inactivated is an "off white" coloured emulsified suspension of highly immunogenic strains of *Pasteurella multocida*, chemically inactivated and oil adjuvanted to get stable emulsion. Selection of highly immunogenic strains of *Pasteurella multocida* for vaccine preparation confers protection to the birds against Fowl Cholera.

The product meets all rigid quality control standards of purity, safety, and potency.

INDICATIONS

This vaccine is recommended for active immunization of birds against Fowl cholera. The vaccine is administered in two doses at the interval of 3 to 4 weeks for effective immune response against this infection.

DOSAGE FORM

Each dose of vaccine contains greater than 10^8 CFU of *Pasteurella multocida* prior to inactivation.

DOSAGE

Dose per bird 0.5 ml

ADMINISTRATION

Shake vaccine bottle well before and during use. Use 20 / 21 gauge needle and automatic syringe, to inject 0.5 ml of vaccine, preferably by subcutaneous route in the lower half of the neck under the loose skin at the point midway between the bird's head and base of the neck, the needle pointing away from the bird's head; or intramuscularly in the breast muscle. Vaccinate using aseptic techniques.

CONTRA-INDICATION

This vaccine is not recommended for those birds, which are clinically sick or severely debilitated, or under conditions of severe stress and strain.

GENERAL RECOMMENDED SCHEDULE OF VACCINATION

Vaccinate layer/broiler breeders and replacement flocks only. Administer the first dose at 8 to 10 weeks age of the bird. Repeat the dose 3 to 4 weeks later, for adequate immune response.

Generally, the second dose is given, a minimum 8 days prior to the predicted age of occurrence of the clinical infection, as per previous experiences of the Veterinarian/Consultant. It is especially recommended to use this vaccine in endemic areas.

SHELF LIFE

Vaccine when stored at recommended conditions has shelf life of 12 months, which is indicated as an expiry date on the label.

STORAGE AND SHIPPING

Vaccine should be stored at +2°C to +8°C in a dark place preferably in the cold room and transported in cold chain system. Do not freeze the vaccine. The pre-chilled vaccine should be transported in thermocol boxes.

PRESENTATION

Bottles of 200 ml and 500 ml capacity, each containing Fowl cholera vaccine of 400 doses and 1000 doses respectively, as specified on the vaccine bottle label.

WARNINGS

- Consult the Veterinarians / Poultry experts regarding vaccination schedule and use of the vaccine.
- This product should be stored, transported, and administered in accordance with the instructions and directions of the manufacturer.
- Bring the vaccine bottle to room temperature.
- Shake vaccine bottle well before and during use.
- Do not use unsterile vaccination equipment.
- Use entire contents when first opened.
- Chemical disinfectants should not be used for sterilization.
- Always vaccinate the healthy birds and keep the vaccinated birds comfortable by providing adequate heat, feed, water, and good ventilation.
- This vaccine is not recommended for the birds that are clinically sick or severely debilitated or under conditions of severe stress and strain.
- If accidentally injected in to human being urgent medical attention is required.
- Burn and destroy the used containers along with the leftover vaccine.



For further information on the vaccine please contact

Venkateshwara Hatcheries Private Limited
(Ventri Biologicals, Vaccine Division)



Mfg. License No. PD-10

Correspondence: 25 km, Pune - Panshet Road, Girinagar, Pune - 411 025, (MS), India.

Plant address: 31 km, Pune - Panshet Road, Village Malkhed, Taluka Haveli, District Pune, (MS), India.

Corporate address: "Venkateshwara House", S. No. 114/A/2, Pune - Sinhagad Road, Pune - 411 030, (MS), India.