



AVIAN INFECTIOUS BRONCHITIS VACCINE, INACTIVATED BP (Vet)

DIRECTIONS FOR USE

DESCRIPTION

Avian Infectious Bronchitis (IB) Vaccine, Inactivated is an "off white" coloured emulsified suspension of highly immunogenic strains of IB virus, chemically inactivated and oil adjuvanted to get stable emulsion. Selection of highly immunogenic strain of IB virus for vaccine preparation confers protection to the birds against IB infection.

The product meets all rigid quality control standards of purity, safety, and potency as per individual monographs of viral vaccines of BP (Vet).

INDICATIONS

This vaccine is recommended for active immunization of birds against Avian Infectious Bronchitis. A dose of vaccine contains high titred inactivated viral antigen, which induce effective immune response against Infectious Bronchitis. The vaccine is recommended for mature birds, which are already primed with IB live vaccine.

DOSAGE FORM

Each dose of vaccine contains greater than 10^6 EID₅₀ of IB virus prior to inactivation.

DOSAGE

Dose per bird 0.5 ml

ADMINISTRATION

Shake vaccine bottle well before and during use. Use 20 / 21 gauge needle and automatic syringe, to inject the vaccine as per dosage, preferably by subcutaneous route in the lower half of the neck under the loose skin at the point midway between the bird's head and base of the neck, the needle pointing away from the bird's head; or intramuscularly in the breast muscle. Vaccinate using aseptic techniques.

CONTRA-INDICATION

This vaccine is not recommended for those birds, which are clinically sick or severely debilitated, or under conditions of severe stress and strain.

GENERAL RECOMMENDED SCHEDULE OF VACCINATION

This vaccine is developed for use in layer and broiler breeder birds. Vaccinate layer and broiler breeder birds between 16 and 20 week of age. The "Midlay" booster dose is recommended for breeder birds under production at the age of 35-40 week for effective protection of progeny of the breeding flock through out the productive life of breeders.

SHELF LIFE

Vaccine when stored at recommended conditions has shelf life of 12 months, which is indicated as an expiry date on the label.

STORAGE AND SHIPPING

Vaccine should be stored at +2°C to +8°C in a dark place preferably in the cold room and transported in cold chain system. Do not freeze the vaccine. The pre-chilled vaccine should be transported in thermocole boxes.

PRESENTATION

Bottles of 200 ml and 500 ml capacity each containing Avian Infectious Bronchitis vaccine of 400 doses and 1000 doses respectively, as specified on the vaccine bottle label.

WARNINGS

- Consult the Veterinarians / Poultry experts regarding vaccination schedule and use of the vaccine.
- This product should be stored, transported, and administered in accordance with the instructions and directions of the manufacturer.
- Bring the vaccine bottle to room temperature.
- Shake vaccine bottle well before and during use.
- Do not use unsterile vaccination equipment.
- Use entire contents when first opened.
- Chemical disinfectants should not be used for sterilization.
- Always vaccinate the healthy birds and keep the vaccinated birds comfortable by providing adequate heat, feed, water, and good ventilation.
- This vaccine is not recommended for the birds that are clinically sick or severely debilitated or under conditions of severe stress and strain.
- If accidentally injected in to human being urgent medical attention is required.
- Burn and destroy the used containers along with the leftover vaccine.



For further information on the vaccine please contact

Venkateshwara Hatcheries Private Limited
(Vetri Biologicals, Vaccine Division)



Mfg. License No. PD-10

Correspondence: 25 km, Pune - Panshet Road, Girinagar, Pune - 411 025, (MS), India.

Plant address: 31 km, Pune - Panshet Road, Village Malkhed, Taluka Haveli, District Pune, (MS), India.

Corporate address: "Venkateshwara House", S. No. 114/A/2, Pune - Sinhagad Road, Pune - 411 030, (MS), India.