



FOWL SPIROCHAETOSIS VACCINE, INACTIVATED

DIRECTIONS FOR USE

DESCRIPTION

Fowl Spirochaetosis Vaccine, Inactivated is an "off white" coloured suspension produced by using local isolates of Spirochaetes grown in Specific Pathogen Free (SPF) chick embryo, chemically inactivated and adjuvanted to get stable emulsion. Selection of highly immunogenic isolate of Spirochaetes for vaccine preparation confers protection to the birds against Spirochaetosis.

The product meets all rigid quality control standards of purity, safety, and potency.

INDICATIONS

This vaccine is used for active immunization of birds against Spirochaetosis.

DOSAGE FORM

Each dose of vaccine contains spirochaete infected embryo mass, as inactivated antigen, 60 - 80 mg per ml prior to inactivation.

DOSAGE

The vaccine is given depending on the body weight of the bird.
Dose per bird 0.5 ml per bird if body weight is less than 1 kg
1.0 ml per bird if body weight is more than 1 kg

ADMINISTRATION

Shake the vaccine bottle well before and during use. Use 20 / 21 gauge needle and automatic syringe, to inject the vaccine intramuscularly using aseptic techniques. Vaccinate the bird depending on the body weight. Administer the vaccine intramuscularly at the rate of 0.5 ml if the weight of the bird is less than 1 kg and if the weight of bird is more than 1 kg inject 1 ml of vaccine.

CONTRA-INDICATION

This vaccine is not recommended for those birds, which are clinically sick or severely debilitated or under conditions of severe stress and strain.

GENERAL RECOMMENDED SCHEDULE OF VACCINATION

It is recommended in the flocks located in Spirochaetosis endemic area. A dose of vaccine contains a large amount of inactivated organisms which act as antigen to induce effective immune response against Spirochaetosis. The Vaccine should be administered atleast 21 days prior to expected outbreak of disease based on previous experience of experts.

SHELF LIFE

Vaccine when stored at recommended conditions has shelf life of 12 months, which is indicated as an expiry date on the label.

STORAGE AND SHIPPING

Vaccine should be stored at +2°C to +8°C in a dark place preferably in the cold room and transported in cold chain system. Do not freeze the vaccine. The pre-chilled vaccine should be transported in thermocole boxes.

PRESENTATION

Amber coloured Bottle of 100 ml capacity containing Spirochaetosis vaccine of 200 / 100 doses as specified on the vaccine bottle label.

WARNINGS

- Consult the Veterinarians / Poultry experts regarding vaccination schedule and use of the vaccine.
- This product should be stored, transported, and administered in accordance with the instructions and directions of the manufacturer.
- Bring the vaccine bottle to room temperature.
- Shake vaccine bottle well before and during use.
- Do not use unsterile vaccination equipment.
- Use entire contents when first opened.
- Chemical disinfectants should not be used for sterilization.
- Always vaccinate the healthy birds and keep the vaccinated birds comfortable by providing adequate heat, feed, water, and good ventilation.
- This vaccine is not recommended for the birds that are clinically sick or severely debilitated or under conditions of severe stress and strain.
- If accidentally injected in to human being urgent medical attention is required.
- Burn and destroy the used containers along with the leftover vaccine.



For further information on the vaccine please contact

Venkateshwara Hatcheries Private Limited
(Ventri Biologicals, Vaccine Division)



Mfg. License No. PD-10

Correspondence: 25 km, Pune - Panshet Road, Girinagar, Pune - 411 025, (MS), India.

Plant address: 31 km, Pune - Panshet Road, Village Malkhed, Taluka Haveli, District Pune, (MS), India.

Corporate address: "Venkateshwara House", S. No. 114/A/2, Pune - Sinhad Road, Pune - 411 030, (MS), India.